

Duality Violation and the $K \rightarrow \pi\pi$ Electroweak Penguin Operator Matrix Elements from Hadronic τ Decays

Kim Maltman^{a,b}, D. Boito^{c,d}, O. Cata^e, M. Golterman^f, M. Jamin^{d,g}, J. Osborne^f, S. Peris^{c,f}

^a*Math & Stats, York U., Toronto, CANADA*

^b*CSSM, U. Adelaide, Adelaide, Australia*

^c*Dept. Physics, U. Autònoma de Barcelona, Barcelona, Spain*

^d*IFAE, U. Autònoma de Barcelona, Barcelona, Spain*

^e*Dept. de Física Teòrica, IFIC, U. Valencia-CSIC, Valencia, Spain*

^f*Dept Physics and Astronomy, SFSU, San Francisco, USA*

^g*ICREA U. Autònoma de Barcelona, Barcelona, Spain*

We discuss a preliminary study of the impact of duality violations on extractions from τ decay data of the $D = 6$ VEVs which determine chiral limit Standard Model $K \rightarrow \pi\pi$ matrix elements of the electroweak penguin operators.

PACS numbers: 12.15.-y, 12.38.Lg, 11.55.Hx, 13.35.Dx

I. INTRODUCTION

In the Standard Model (SM), ϵ'/ϵ is dominated by contributions from the gluonic and electroweak penguin (EWP) operators, Q_6 and Q_8 . In the $SU(3)$ chiral limit, the $K \rightarrow \pi\pi$ matrix elements of the EWP operators $Q_{7,8}$, are determined by two 4-quark VEVs, $\langle \mathcal{O}_1 \rangle$ and $\langle \mathcal{O}_8 \rangle$, which also determine the dimension $D = 6$ part of the OPE of the flavor ud V-A correlator difference $\Delta\Pi \equiv \Pi_V^{(0+1)} - \Pi_A^{(0+1)}$ [1], where the superscript $0+1$ denotes the sum of spin $J = 0$ and 1 components and, with $J_{V/A}^\mu$ the standard V or A ud current, the scalar correlators $\Pi_{V/A}^{(J)}$ are defined via


$$i \int d^4x e^{iq \cdot x} \langle 0 | T(J_{V/A}^\mu(x) J_{V/A}^\nu(0)^\dagger) | 0 \rangle \equiv (q^\mu q^\nu - g^{\mu\nu} q^2) \Pi_{V/A}^{(1)}(q^2) + q^\mu q^\nu \Pi_{V/A}^{(0)}(q^2). \quad (1)$$

Since $[\Delta\Pi]_{D=6}^{OPE}$ is strongly dominated by the contribution involving $\langle \mathcal{O}_8 \rangle$, which VEV also dominates the chiral limit Q_8 matrix element, the extraction of $[\Delta\Pi]_{D=6}^{OPE}$ is of considerable phenomenological interest, and a number of dispersive and finite energy sum rule (FESR) analyses have attempted it [2–5]. τ decay data plays a key role in these analyses since the spectral function of $\Delta\Pi$, $\Delta\rho(s) = \frac{1}{\pi} \text{Im} \Delta\Pi(s+i\epsilon)$, is directly measurable for $s \leq m_\tau^2$ in non-strange hadronic τ decays. Explicitly, in the SM, with S_{EW} a short-distance EW correction, $y \equiv s/m_\tau^2$, $w_T(y) = (1-y)^2(1+2y)$ and $R_{V/A}^{ud} \equiv \Gamma[\tau^- \rightarrow \nu_\tau \text{ hadrons}_{V/A}^{ud}]/\Gamma[\tau^- \rightarrow \nu_\tau e^- \bar{\nu}_e]$, one has, for the continuum (non- π -pole) part of $\Delta\rho$ [6]

$$\Delta\rho(s) = \frac{m_\tau^2}{12\pi^2 |V_{ud}|^2 S_{EW} w_T(y)} \frac{dR_{V-A}^{ud}}{ds}. \quad (2)$$

Dispersive analyses employ the unsubtracted dispersion relation for $\Delta\Pi$ and require either assumptions about the saturation of the dispersion integral within the range kinematically accessible in τ decays, or supplementary constraints on $\Delta\rho(s)$ for $s > m_\tau^2$, such as those

FIG. 1: Residuals for the CGM “maximally safe” fit, with no DVs



provided by the Weinberg sum rules [7] and the DGMLY π electromagnetic (EM) self-energy sum rule [8] (see, e.g., Ref. [2] for details). Higher dimension ($D > 6$) contributions to $\Delta\Pi(Q^2)$ must also be considered. These problems are avoided in the FESR approach, which relies on $\Delta\Pi(s)$ having no kinematic singularities and hence satisfying the FESR relation


$$\int ds w(s) \Delta\rho(s) = \frac{-1}{2\pi i} \oint_{|s|=s_0} ds w(s) \Delta\Pi(s), \quad (3)$$

for any s_0 and any $w(s)$ analytic in the region of the contour. For sufficiently large s_0 , the OPE should become reliable on the RHS. Choosing polynomial weights $w(s)$ with degree N strongly suppresses OPE contributions with $D > 2N + 2$. For sub-asymptotic s_0 , OPE breakdown, or duality violation (DV) is expected. In fact, even for $s_0 \sim m_\tau^2$, sizeable s_0 -dependent deviations between the LHS and OPE versions of the RHS are found for the $w(s) = 1$ V and A analogues of Eq. 3 [9, 10]. These are strongly suppressed for analogues employing pinched weights ($w(s)$ with a zero at $s = s_0$) [9], indicating that at scales $\sim m_\tau^2$ DVs are localized to the vicinity of the timelike axis. With this in mind, the analysis of Ref. [4] (CGM) employed doubly pinched weights, checking the s_0 -dependence of the match between the weighted spectral integrals and optimized OPE fit as a test of the self-consistency of the assumed neglect of residual DV contributions. Figure 1 shows the resulting residuals, $[I_{OPE}^w(s_0) - I_{OPAL}^w(s_0)] / \delta I_{OPAL}^w(s_0)$, over an expanded s_0 range, for the two weights, w_1 and w_2 of the “maximally safe” CGM analysis based on OPAL data [11]. (We focus here on OPAL data due to a problem with the ALEPH covariance matrices [10] which is the subject of ongoing reanalysis.) $I_{OPAL, OPE}^w(s_0)$ are the LHS and RHSs of Eq. 3 and $\delta I_{OPAL}^w(s_0)$ the uncertainty on $I_{OPAL}^w(s_0)$. It is obvious that residual DVs, though not evident *within errors* above $s_0 \sim 2 \text{ GeV}^2$, become non-negligible below this point. Small residual DV contributions are thus expected in the $s_0 > 2 \text{ GeV}^2$ CGM fit window as well. Lacking a model for DVs, analyses such as CGM were unable to estimate the systematic uncertainty associated with neglecting these contributions.

II. INCORPORATING DUALITY VIOLATIONS

In Refs. [12], a model for DV spectral contributions was developed. The model builds on earlier work in Refs. [13] and is motivated by large- N_c and Regge-based resonance spacing

FIG. 2: The Weinberg and DGMLY sum rule tests, the DGMLY case involving the weight $w(s) = s \log(s/\Lambda^2)$.



ideas. The model leads to ansätze $\rho_{V/A}(s) = \rho_{OPE}^{(D=0)}(s) + \rho_{V/A}^{DV}(s)$, $s > s_{min}$, for the V and A spectral functions, where the DV contributions have the form

$$\rho_{V/A}^{DV}(s) = \kappa_{V/A} e^{-\gamma_{V/A}s} \sin(\alpha_{V/A} + \beta_{V/A}s). \quad (4)$$

In Refs. [5] the impact of DVs on previous V-A analyses was investigated using a *single* DV ansatz of the form Eq. 4 for the V-A difference $\Delta\rho(s)$. This involves the implicit additional assumption that $\beta_V \simeq \beta_A$ and $\gamma_V \simeq \gamma_A$, allowing the 8-parameter V-A difference to be re-written in the effective 4-parameter form, Eq. 4. We avoid this additional assumption and fit the V and A DV parameter sets separately, as part of a combined V, A fit which also determines the OPE parameters α_s , $\langle\alpha_s G^2\rangle$, and the relevant $D = 6$ and 8 V and A channel effective condensates. We find central DV parameter fit values not in good accord with the expectations $\beta_V \simeq \beta_A$, $\gamma_V \simeq \gamma_A$.

Our analysis employs $w(s)$ up to degree 3, including $w(s) = 1$, which is optimally sensitive to the DV contributions. The resulting fits provide excellent matches between the OPAL spectral integrals and optimized OPE+DV fit forms for all $w(s)$ employed and all s_0 down to a fit window minimum $s_0^{min} \sim 1.4 - 1.5 \text{ GeV}^2$. Though so far aimed at extracting α_s , and not optimized for extracting $D = 6, 8$ V-A condensates, the analysis nonetheless provides preliminary results for these quantities. Since the fits provide a prediction for $\Delta\rho(s)$ for $s > s_0^{min}$, and hence also above $s = m_\tau^2$, we can test our results against the Weinberg and DGMLY sum rules, which constraints have *not* been incorporated in performing the fits. The first and second Weinberg sum rules are written in a form with RHSs equal to zero; for the RHS of the DGMLY sum rule we employ the $SU(2)$ chiral limit value $-8\pi f_\pi^2 [\delta m_\pi^2]_{EM} / 3\alpha_{EM} = -0.0109(15) \text{ GeV}^4$ [5]. The results of these tests are shown in Figure 2, with s_0 the point beyond which the fitted form of $\Delta\rho(s)$ is employed in the relevant spectral integral. Below this point, experimental data are used. The dotted and solid black lines in the third panel show the central DGMLY sum rule RHS and error. All three sum rules should be satisfied for all s_0 in our fit window. This is evidently the case, giving us good confidence in the results for the fitted OPE parameters as well.

As an illustration of our preliminary results, we quote the effective $D = 6$ V-A condensate C_6^{V-A} which results from an update of the CGM OPAL fit incorporating DVs and fitting ρ_V , ρ_A , and the $w(y) = 1$, $1 - y$ and $(1 - y)^2$ ($y = s/s_0$) FESRs with the CIPT scheme for the truncated $D = 0$ OPE series and $s_0^{min} = 1.504 \text{ GeV}^2$. C_6^{V-A} is defined by $[\Delta\Pi(Q^2)]_{D=6}^{OPE} = C_6^{V-A}/Q^6$. We find $C_6^{V-A} = -0.0057(18) \text{ GeV}^6$, c.f. the CGM no-DVs maximally safe

analysis OPAL-based value $-0.0054(8)$ GeV⁶. Values somewhat larger in magnitude, with larger errors, are obtained from analyses excluding $w(y) = 1 - y$, including those employing FOPT for the $D = 0$ series. We emphasize that these results are preliminary, and that a dedicated V-A analysis, aimed at reducing the errors, is in progress. Readers noting the 20 – 25% difference between the values quoted above and those obtained from the ALEPH-based analyses of CGM (neglecting DVs) and Ref. [5] (including DVs approximately) should bear in mind that the ALEPH and OPAL 4π data differ significantly in the upper part of the spectrum, with the OPAL data agreeing better with expectations based on CVC and recent preliminary BaBar and SND 4π electroproduction data [14].

Acknowledgments

This work was supported by the Natural Sciences and Engineering Research Council of Canada, the US DOE, MICINN (Spain) under Grant FPA 2007-60323, the Spanish Consolider Ingenio 2010 Program CPAN (CSD2007-00042), CICYTFEDER-FPA2008-01430, SGR2005-00916 and the Programa de Movilidad PR2010-0284.

-
- [1] J.F. Donoghue and E. Golowich, *Phys. Lett.* **478**, 172 (2000).
 - [2] V. Cirigliano *et al.*, *Phys. Lett.* **B522**, 245 (2001).
 - [3] S. Peris, B. Phily, E. de Rafael, *Phys. Rev. Lett.* **86**, 14 (2001); J. Bijnens, E. Gamiz, J. Prades, *JHEP* **0110**, 009 (2001); M. Knecht, S. Peris, E. de Rafael, *Phys. Lett.* **B508**, 117 (2001); B.L. Ioffe, K.N. Zyablyuk, *Nucl. Phys.* **A687**, 437 (2001); K. N. Zyablyuk, *Eur. Phys. J.* **C38**, 215 (2004); J. Roja, J.I. Latorre, *JHEP* **0401**, 055 (2004); C. A. Dominguez, K. Schilcher, *Phys. Lett.* **B581**, 193 (2004); S. Friot, D. Greynat, E. de Rafael, *JHEP* **0410**, 043 (2004); S. Narison, *Phys. Lett.* **B624**, 223 (2005); S. Schael *et al.* (ALEPH), *Phys. Rep.* **421**, 191 (2005); J. Bordes *et al.*, *JHEP* **0602**, 037 (2006); P. Masjuan, S. Peris, *JHEP* **0705**, 040 (2007); A.A. Almasly, K. Schilcher, H. Spiesberger, *Eur. Phys. J.* **C55**, 237 (2008).
 - [4] V. Cirigliano *et al.*, *Phys. Lett.* **B555**, 71 (2003); V. Cirigliano, E. Golowich and K. Maltman, *Phys. Rev.* **D68**, 054013 (2003).
 - [5] M. Gonzalez-Alonso, A. Pich and J. Prades, *Phys. Rev.* **D81**, 074007 (2010); *Phys. Rev.* **D82**, 014019 (2010).
 - [6] Y. Tsai, *Phys. Rev.* **D4**, 2821 (1971).
 - [7] S. Weinberg, *Phys. Rev. Lett.* **18**, 507 (1967).
 - [8] T. Das *et al.*, *Phys. Rev. Lett.* **18**, 759 (1967).
 - [9] K. Maltman, *Phys. Lett.* **B440**, 367 (1998); C. A. Dominguez and K. Schilcher, *Phys. Lett.* **448**, 93 (1999).
 - [10] D. Boito *et al.*, arXiv: 1011.4426 [hep-ph]; arXiv: 1103.4194 [hep-ph]
 - [11] K. Ackerstaff *et al.* (OPAL), *Eur. Phys. J.* **C7**, 571 (1999).
 - [12] O. Catà, M. Golterman and S. Peris, *JHEP* **0508**, 076 (2005); *Phys. Rev.* **D77**, 093006 (2008); *Phys. Rev.* **D79**, 053002 (2009).
 - [13] B. Blok, M.A. Shifman and D.X. Zhang, *Phys. Rev.* **D57**, 2691 (1998) [erratum: *ibid.* 19901 (1999)], I.I.Y. Bigi, *et al.*, *Phys. Rev.* **D59**, 054011 (1999).
 - [14] See, e.g., V.P. Druzhinin *et al.*, arXiv:1105.4975 and the poster session presentation by L. Kardapoltsev at PANIC 2011.